

PolyVision Brings You Good Vision and Healthy Eyes

看得清楚嗎? See this clearly


距離4米,遮左/右眼測試。視力表 只作參考,並不等同至面眼科視光檢查。

Cover your left / right eye and read the letter at 4 meters. This simple test is not equivalent to a comprehensive eye examination.



為什麼您應該定期接受「全面眼科視光檢查」?

眼睛是我們的靈魂之窗,隨著年紀增長,雙眼亦會不斷變化,除了屈光度數上的改變,我們更關心眼睛的內外健康。 Vision is one of our most important sensory functions. Just like any other organs, the condition of our eyes changes with time.



好好保護眼睛

光線充足 Sufficient Lighting

















正確坐姿 Good Posture





常見症狀Check—Check Common Symptoms Checklist

- 眼攰、眼乾、眼澀 Tired, dry, gritty eyes
- ② 視力糢糊或時矇時清 Blurry / fluctuating vision ⑦ 過近距離閱讀/看電視 Close reading / TV watching distance
- **3** 對焦困難 Difficult to focus
- 4 重影 Double vision
- 6 畏光、對光線敏感 Sensitive to light
- 6 經常捽眼、流眼水 Rubbing eyes, tearing frequently
- ⑧ 矇眼、單眼或側頭 Squinting, covering one eye or head tilting to one side
- 眨眼過度頻密 Excessive eye blinking
- ⑩ 閱讀時跳行跳字 Losing place when reading

了解您的眼睛 Learn About Your Eye

近視 Myopia (near/short-sightedness)

- 因眼球過長,影像對焦在視網膜前面,看遠景物糢糊不清 Eyeball is too long, visual image is formed in front of retina and causes blurry distance vision.
- 分先天遺傳、種族和後天用眼習慣、環境等 Partly inherited and partly due to environmental factors.

(遠視 Hyperopia (far/long-sightedness)


- 因眼球過短,影像對焦在視網膜後面,視乎度數及眼睛調節力,看近(甚至 看遠)景物糢糊不清
- Eyeball is too short, visual image is formed behind retina. Depends on person's prescription and accommodative function, it may lead to blurry vision at near or even distance.
- 多屬先天性,有機會隨年齡減淺 Usually congenital and may decrease with age.

(斜視 Strabismus (Squint))

- 因眼外肌不協調或雙眼度數偏差過大
- Caused by poor coordination of eye muscles or significant difference in refractive error.
- 視軸不平衡,雙眼望向不同方向
 Eyes are misaligned and point towards different directions.
- 分內/外、上/下、顯性/隱性斜視 Classified into esotropia / exotropia, hypertropia / hypotropia, manifest or latent squint.

弱視 Amblyopia (懶惰眼 lazy eye)

- 因深度數、雙眼度數偏差過大或斜視 Caused by high or significant difference in refractive errors or strabismus.
- 眼睛未能提供清晰影像,刺激腦部視覺發展或清晰影像被抑制
 No clear image to stimulate visual brain development or clear image is suppressed.
- 即使配戴正確度數的眼鏡,視物仍糢糊不清 Cannot achieve clear vision even with aid of glasses
- 多出現在一邊眼睛,不易察覺,延誤處理可引致永久弱視
 Normally affect one eye only, often goes undetected and delay in treatment may lead to permanent impairment.



全面眼科視光檢查 Comprehensive Eye Examination 青光眼檢查 Glaucoma Examination 糖尿眼檢查 Diabetes Eye Examination 角膜矯形 Orthokeratology

學習視覺檢查 School Vision Assessment 兒童視覺訓練 Children Vision Training 色覺檢查 Color Vision Testing 視野檢查 Visual Field Testing 眼球結構斷層掃描 Cross-sectional Ocular Scan (Optical Coherence Tomography) 可溶性淚管塞 Dissolvable Punctal Plug Occlusion 隱形眼鏡驗配及跟進護理 Contact Lens Fitting and Aftercare 眼鏡及太陽眼鏡驗配 Optical Dispensing of Spectacles and Sunglasses

飛蚊症 Floaters

or detachment.

白內障 Cataract

因玻璃體變得混濁或有飄浮物

脱落等嚴重眼疾的先兆

• 晶狀體變得混濁、不透明

• 因年齡增長、遺傳、吸煙等

blindness.

青光眼 Glaucoma

eventually blindness.

Small floating debris in vitreous humour.

因年齡增長、紫外線、疾病、創傷等

【 黃斑病變 Macular Degeneration 】

• 黄斑逐漸退化、出現滲漏、出血

Cause by ageing, UV, diseases, injury, etc.

Lens becomes cloudy and not transparent.

• 視覺逐漸糢糊、出現光圈、顏色暗淡、畏光等

Associated with aging, family history, smoking etc.

分急性(視力突減、眼紅、眼痛)、慢性(病徵不明顯)

• 視神經受損、視野收窄、視力下降,嚴重可致盲

Optic nerve damage, visual field and vision loss, or

chronic glaucoma (no obvious symptoms).

• 不同形狀或大小的黑點或暗影在眼前浮動

Different shapes and sizes, e.g. tiny shadowy dots or strands.

• 看見大量「飛蚊」或持續性「閃光」, 亦可能是視網膜裂孔

Blurry vision, glare, reduced color perception, sensitive to light, etc.

Gradual degeneration of macular region, blood vessels leakage or bleeding.

Blurry and distorted central vision, sudden vision loss, or eventually

Classified into acute glaucoma (sudden vision loss, red and painful eye) or

• 中央視力糢糊、彎曲、變形、暗點、視力驟降,嚴重可致盲

Sudden increase of floaters and/or flashes may be sign of retinal breaks





理大護眼(銅鑼灣)中心 PolyVision Eyecare (Causeway Bay) Centre

香港銅鑼灣勿地臣街1號 時代廣場1座1703-1705室 Room 1703-1705, Tower One, Times Square, 1 Matheson Street, Causeway Bay, Hong Kong 電話 Tel: 2866 9660

理大護眼(太古)中心_____

PolyVision Eyecare (Taikoo) Centre

香港鰂魚涌康山道1號 康怡廣場辦公大樓1201-1203室 Room 1201-1203, Kornhill Plaza Office Tower, 1 Kornhill Road, Quarry Bay, Hong Kong 電話 Tel: 2377 9660

理大護眼(尖沙咀)中心 PolyVision Eyecare (Tsimshatsui) Centre

九龍尖沙咀廣東道5號 海港城海洋中心1217-1218室 Room 1217-1218, Ocean Centre, Harbour City, 5 Canton Road, Tsim Sha Tsui, Kowloon 電話 Tel: 2750 9660

理大護眼(旺角)中心 PolyVision Eyecare (Mongkok) Centre

九龍旺角彌敦道639號 雅蘭中心辦公樓1期2102-2105室 Room 2102-2105, Office Tower One, Grand Plaza, 639 Nathan Road, Mongkok, Kowloon 電話 Tel: 2780 9660

理大護眼(沙田)中心 PolyVision Eyecare (Shatin) Centre

新界沙田白鶴汀街10-18號 新城市商業大廈810-812室 Room 810-812, New Town Tower, 10-18 Pak Hok Ting Street, Shatin, New Territories 電話 Tel: 2153 9660

理大護眼(荃灣)中心 PolyVision Eyecare (Tsuen Wan) Centre

新界荃灣眾安街68號 荃灣千色匯1期2301-02及07-08室 Room 2301-02 & 07-08, KOLOUR • Tsuen Wan I, 68 Chung On Street, Tsuen Wan, New Territories 電話 Tel: 2245 9660

服務時間 Service Hours

 星期一至五 Monday - Friday
 10:00 - 19:00


 星期六 Saturday
 09:00 - 18:00

 星期日 Sunday
 14:00 - 18:00

 公眾假期休息 Closed on Public Holiday

www.polyvision.com.hk

黃斑病變阿姆斯勒柵絡檢查 | Amsler Grid Test for Macular Degeneration |





距離30厘米,遮左/右眼,凝視中心白點(如有需要請配戴閱讀眼鏡)。 感覺方格線條有否出現扭曲、斷裂、 變形、糢糊、暗點或空缺。 Cover your left / right eye and hold this at 30cm distance from your eyes. While

30cm distance from your eyes. While focusing on the central dot, look for any wavy, blurry lines and check for missing areas, lines and corners.

(Wear reading glasses if needed.)



→站式優質護眼服務 One-Stop Quality Eyecare Service